

## AMENDMENTS TO THE CLAIMS

### Listing of Claims:

The following list of claims replaces all previous listings or versions thereof:

1. (Currently amended) A ~~highly-sensitive-real-time RT-PCR capable of~~method for specifically detecting the expression of more than one MAGE gene, ~~wherein comprising~~ reverse transcription of ~~the corresponding~~ MAGE transcripts ~~is carried out~~ simultaneously performed in a single cDNA-synthesis reaction with a cDNA-primer MgRT3a consisting of SEQ ID NO:4.
- 2-18. (Canceled)
19. (Currently amended) A diagnostic composition comprising a cDNA-primer MgRT3a consisting of [[O]SEQ ID NO:4[D]].
20. (Canceled)
21. (Previously presented) An oligonucleotide designated MgRT3a (SEQ ID NO:4).
- 22-23. (Canceled)
24. (Currently amended) The diagnostic composition of claim 19, further comprising one or more of the primerprimers Mg1\_RT5a (SEQ ID NO:14), MgRT2 (SEQ ID NO:XX), MgRT1b (SEQ ID NO:XX), MgRT4 (SEQ ID NO:XX), MgRT6 (SEQ ID NO:XX), MgRT1a (SEQ ID NO:XX), MgRT3b (SEQ ID NO:XX), MgRT5b (SEQ ID NO:XX), Mg1\_RT1 (SEQ ID NO:XX), Mg1\_RT2 (SEQ ID NO:XX), Mg1\_RT3 (SEQ ID NO:XX), Mg1\_RT4 (SEQ ID NO:XX), Mg1\_RT5c (SEQ ID NO:XX), Mg1\_RT5d (SEQ ID NO:XX), Mg1\_RT5e (SEQ ID NO:XX), Mg1\_RT6 (SEQ ID NO:XX), and Mg1\_RT7 (SEQ ID NO:XX).
- 25-26. (Canceled)

27. (Currently amended) The diagnostic composition of claim 19, the composition further comprising a cDNA-primer that hybridizes to a calibrator mRNA for reverse transcription of a calibrator mRNA.
28. (Currently amended) The diagnostic composition of claim 27, wherein the calibrator mRNA is porphobilinogen desaminase (PBGD) mRNA.
29. (Canceled)
30. (Currently amended) The diagnostic composition of claim 28, wherein the composition ~~further comprises oligonucleotide MgRT3a (SEQ ID NO:4) as a primer for reverse transcription of the at least two different MAGE gene transcripts and PBGD RT15b (SEQ ID NO: 35) as primer for reverse transcription of the PBGD mRNA.~~
31. (Currently amended) The diagnostic composition of claim 28, ~~the composition further comprising PCR primers for amplification of the calibrator mRNA, wherein the calibrator mRNA is porphobilinogen desaminase (PBGD) mRNA, and wherein said PCR-primers for amplification of PBGD-cDNA comprise the oligonucleotides hu PBGD se (SEQ ID NO:44) and PBGD R (SEQ ID NO:50) as primer pairs for PCR amplification of PBGD-cDNA.~~
32. (Canceled)
33. (Currently amended) The diagnostic composition of claim 19, ~~the composition further comprising PCR-primers for amplification of MAGE-cDNA, the primers comprising oligonucleotides selected from one of the following groups:~~

(C)

PCR-primer	sequence (5' - 3')
<u>MAGE-A1</u>	<u>GTA GAG TTC GGC CGA AGG AAC</u>
<u>MAGE-A1</u>	<u>CAG GAG CTG GGC AAT GAA GAC</u>
<u>MAGE-A2</u>	<u>CAT TGA AGG AGA AGA TCT GCC T</u>
<u>MAGE-A2</u>	<u>GAG TAG AAG AGG AAG AAG CGG T</u>

<u>MAGE-A3/6</u>	<u>GAA GCC GGC CCA GGC TCG</u>
<u>MAGE-A3/6</u>	<u>GAT GAC TCT GGT CAG GGC AA</u>
<u>MAGE-A4</u>	<u>CAC CAA GGA GAA GAT CTG CCT</u>
<u>MAGE-A4</u>	<u>TCC TCA GTA GTA GGA GCC TGT</u>
<u>MAGE-A10</u>	<u>CTA CAG ACA CAG TGG GTC GC</u>
<u>MAGE-A10</u>	<u>GCT TGG TAT TAG AGG ATA GCA G</u>
<u>MAGE-A12</u>	<u>TCC GTG AGG AGG CAA GGT TC</u>
<u>MAGE-A12</u>	<u>ATC GGA TTG ACT CCA GAG AGT A</u>

(D)

<b>PCR-primer</b>	<b>sequence (5' - 3')</b>
<u>MAGE-A1</u>	<u>TAG AGT TCG GCC GAA GGA AC</u>
<u>MAGE-A1</u>	<u>CTG GGC AAT GAA GAC CCA CA</u>
<u>MAGE-A2</u>	<u>CAT TGA AGG AGA AGA TCT GCC T</u>
<u>MAGE-A2</u>	<u>CAG GCT TGC AGT GCT GAC TC</u>
<u>MAGE-A3/6</u>	<u>GGC TCG GTG AGG AGG CAA G</u>
<u>MAGE-A3/6</u>	<u>GAT GAC TCT GGT CAG GGC AA</u>
<u>MAGE-A4</u>	<u>CAC CAA GGA GAA GAT CTG CCT</u>
<u>MAGE-A4</u>	<u>CAG GCT TGC AGT GCT GAC TCT</u>
<u>MAGE-A10</u>	<u>ATC TGA CAA GAG TCC AGG TTC</u>
<u>MAGE-A10</u>	<u>CGC TGA CGC TTT GGA GCT C</u>
<u>MAGE-A12</u>	<u>TCC GTG AGG AGG CAA GGT TC</u>
<u>MAGE-A12</u>	<u>GAG CCT GCG CAC CCA CCA A</u>

34. (Canceled)